

Ecosystems (Unity and diversity) revision notes

This group covers how new species arise and how biodiversity is being lost. A4.1 sets out the evidence for evolution and the process of speciation. A4.2 covers biodiversity, the causes of the current extinction crisis and the approaches used to conserve it.

For IB Diploma Biology SL undefined. **HL only** is additional higher level content, for HL students only.

1 Evolution and speciation Spec A4.1

Evolution as change in heritable characteristics Spec A4.1.1

Evolution is change in the **heritable characteristics** of a **population**. Heritable means coded in the genes and passed from parents to offspring. Evolution therefore happens to populations over generations, not to individuals during their lives.

At the level of genes, evolution is a change in the frequency of alleles in the gene pool of a population. A new allele may arise by mutation, then become more or less common in later generations.

A characteristic an organism acquires during its life is not evolution. Larger muscles from exercise, a scar, or a plant growing tall in fertile soil do not change the DNA in gametes, so they are not inherited.

Darwinian evolution and Lamarckism compared		
	Lamarckism	Darwinian evolution by natural selection
Where variation comes from	Organisms change during their lifetime through use or disuse of structures	Heritable variation already exists between individuals, from mutation and sexual reproduction
What is inherited	Acquired characteristics	Only genetic (heritable) characteristics
What changes	Individuals	The population: allele frequencies shift over generations
Status	Not supported: acquired changes do not alter the genes passed on in gametes	Supported by a wide range of independent evidence

Common mistake: Do not write that an individual organism evolves or adapts in order to survive. Individuals survive or die; the population evolves.

Exam tip: Nature of science: evolution by natural selection is called a theory because no theory can be formally proved true, not because it is weak. It predicts and explains a broad range of observations and is unlikely ever to be falsified.

Evidence from sequences and selective breeding Spec A4.1.2, A4.1.3

Base and amino acid sequences

All cellular organisms store genetic information as DNA and use almost the same genetic code. This shared chemistry points to a single common origin of life. Comparing sequences between species gives **powerful evidence of common ancestry**.


- Mutations change base sequences over time. Once two populations have separated, each lineage accumulates its own mutations.
- So the **fewer differences** between the base sequences of a gene (or the amino acid sequences of a protein) in two species, the **more recently** they shared a common ancestor.
- Proteins found in a very wide range of species, such as cytochrome c (used in respiration) and haemoglobin, are often compared.
- If differences accumulate at a roughly steady rate, the number of differences can also be used to estimate how long ago two lineages diverged.

Base sequences show more differences than amino acid sequences. The genetic code is degenerate, so some base substitutions do not change the amino acid.

Sequence evidence is independent of anatomy, yet the relationships it shows mostly agree with those based on structure. When independent lines of evidence point the same way, the evidence is strong.

position	1	2	3	4	5	6	7	8	9	10	11	12	differences from human
human	Gly	Asp	Val	Glu	Lys	Gly	Lys	Lys	Ile	Phe	Ile	Met	—
species X	Gly	Asp	Ile	Glu	Lys	Gly	Lys	Lys	Ile	Leu	Ile	Met	2
species Y	Ala	Asp	Thr	Glu	Lys	Ala	Lys	Arg	Ile	Tyr	Ile	Leu	6

 = amino acid differs from the human sequence (illustrative sequences)



fewer differences = more recent common ancestor

[\(opens full size in a new tab\)](#)

==Fewer differences = more recent common ancestor==: human and X differ at 2 positions, Y at 6 (illustrative sequences).

Selective breeding

In **selective breeding** (artificial selection), humans choose individuals with desired traits to breed, generation after generation. Domesticated breeds and crop varieties now differ greatly from each other and from the original wild species.

- Wild cabbage (*Brassica oleracea*) has been bred into cabbage, kale, broccoli, cauliflower, Brussels sprouts and kohlrabi, by selecting for large leaves, flower buds, side buds or swollen stems.
- All dog breeds, from chihuahuas to great Danes, descend from domesticated wolves.
- Maize was bred from the wild grass teosinte, which has tiny ears with a few hard-cased grains.

This shows how **rapidly** evolutionary change can occur: large heritable changes in a few thousand years or less. If human selection can do this, natural selection acting over millions of years can produce far greater changes.

Common mistake: Selective breeding is evidence that populations can change, but it is not natural selection. The selecting agent is humans, not the environment.


Homologous and analogous structures Spec A4.1.4, A4.1.5

Homologous structures have the same basic structure in different species because they were inherited from a **common ancestor**, even though they may now have different functions. They are evidence of common ancestry followed by divergence.

The pentadactyl limb

The limbs of amphibians, reptiles, birds and mammals are built on the same five-digit (**pentadactyl**) plan:

- one bone in the upper limb: humerus (forelimb) or femur (hindlimb)
- two bones in the lower limb: radius and ulna, or tibia and fibula
- a group of wrist or ankle bones: carpals or tarsals
- a set of bones in the palm or sole: metacarpals or metatarsals
- five digits made of phalanges



[\(opens full size in a new tab\)](#)

The same bones in a human arm, bat wing and whale flipper: humerus, radius and ulna, carpals (wrist), metacarpals (hand) and phalanges (fingers).

One plan, different functions		
Forelimb	Function	How the plan is modified
Human arm	Grasping and manipulating	Long separate digits with an opposable thumb
Bat wing	Flying	Very long metacarpals and phalanges support a membrane of skin
Whale flipper	Swimming	Short, flattened bones enclosed in a paddle
Bird wing	Flying	Carpals and metacarpals fused, digits reduced

There is no functional reason why a flipper, a wing and a hand need the same bones. The shared plan is best explained by descent from a common ancestor with a pentadactyl limb.

Convergent evolution and analogous structures

Analogous structures have the same function but different evolutionary origins. They arise by **convergent evolution**: unrelated lineages meet similar selection pressures, so natural selection produces similar adaptations independently.

- Wings of birds and insects: a bird wing is a modified forelimb with bones and feathers; an insect wing is an outgrowth of the exoskeleton with no bones.
- Camera eyes of octopuses and vertebrates: both focus light with a lens onto a retina, but they develop differently and the octopus retina has no blind spot.
- Streamlined bodies and fins of sharks (fish) and dolphins (mammals).

Homologous and analogous structures		
	Homologous	Analogous
Evolutionary origin	Same, from a common ancestor	Different
Function	Often different	Same
Type of evolution	Divergent	Convergent
Example	Pentadactyl limbs	Bird and insect wings

Exam tip: Linking question: what counts as strong evidence in biology? Homologous structures, sequence data and selective breeding are independent lines of evidence that all support common ancestry.

Speciation, reproductive isolation and differential selection Spec A4.1.6, A4.1.7

Speciation is the formation of new species by the **splitting of a pre-existing species**. One population divides into two or more groups that evolve separately until they can no longer interbreed to produce fertile offspring. This is the only way in which new species have appeared.

- Speciation increases the total number of species on Earth. Extinction decreases it.
- Gradual evolutionary change within a single species over time is not speciation, because the number of species does not increase.

How a species splits

1. **Reproductive isolation**: two populations of one species stop interbreeding, so there is no gene flow between them.
2. **Differential selection**: the populations live under different conditions, so natural selection favours different alleles in each.
3. Different mutations also arise in each population, and chance changes in allele frequency happen separately.


4. Over many generations the gene pools diverge. Eventually the populations could not interbreed and produce fertile offspring even if they met. They are now separate species.

Geographical isolation is one way of achieving reproductive isolation: a physical barrier such as a river, a mountain range or the sea separates the populations.

Bonobos and common chimpanzees

Bonobos (*Pan paniscus*) and common chimpanzees (*Pan troglodytes*) are closely related species in the genus *Pan*. Genetic estimates put the split between the two species at roughly 1 to 2 million years ago. The traditional explanation is that the Congo River formed at about this time and divided the ancestral population. Geological evidence suggests the river is much older, so another hypothesis is that ancestral bonobos crossed to the south bank during a period of unusually low river flow and were then cut off. Either way, the river is the barrier: these apes do not swim, so there is no gene flow across it. Bonobos live only south of the river; common chimpanzees live to the north.

The two populations then faced different selection pressures. One widely discussed explanation is that bonobos, south of the river, had no gorillas competing for ground vegetation, so food was more reliable. This favoured large, tolerant, female-led groups with little aggression. North of the river, chimpanzees share their range with gorillas. This favoured male-dominated groups, more aggression and cooperative hunting.



[\(opens full size in a new tab\)](#)

The Congo River is a **geographical barrier**: no gene flow, different selection pressures on each bank.

Exam tip: In speciation answers name both stages: reproductive isolation stops gene flow, and differential selection makes the isolated gene pools diverge. Isolation alone does not create a new species.

Quick check

- 1 . Define evolution.
- 2 . What do fewer differences in the amino acid sequence of a protein between two species suggest?
- 3 . Name the five groups of bones in a pentadactyl forelimb.
- 4 . What two processes are needed for speciation?

2 Conservation of biodiversity Spec A4.2

Biodiversity and the number of species Spec A4.2.1, A4.2.2

Biodiversity is the variety of life in all its forms, levels and combinations. It is usually considered at three levels.

Three levels of biodiversity		
Level	Meaning	Example
Ecosystem diversity	The variety of habitats and ecosystems in an area	A coastline with coral reefs, seagrass beds, mangroves and desert
Species diversity	The number of different species (richness) and how evenly individuals are spread between them (evenness)	A tropical rainforest has more tree species than a pine plantation
Genetic diversity	The variety of alleles within a species or population	The many varieties of rice; the low genetic diversity of cheetahs

Genetic diversity matters because a population with more alleles is more likely to include individuals that can survive a new disease or a change in conditions.

How many species are there?

About two million species have been **discovered, named and described**, but there are **many more species to be discovered**. Most estimates of the total run to several million eukaryote species, and most bacteria and archaea have not been described. Many undescribed species are insects, fungi and organisms of tropical forests and the deep sea.

Evidence from fossils suggests that there are **more species alive on Earth today than at any time in the past**. Over long periods speciation has outpaced extinction, despite five earlier mass extinctions. The fossil record is incomplete, so past numbers are estimates.

Exam tip: Nature of science: classification is pattern recognition, and the same observations can be classified differently. Splitters recognise more species than lumpers in a taxonomic group, so species counts depend partly on who does the counting.

Anthropogenic species extinction and ecosystem loss Spec A4.2.3, A4.2.4

The current **sixth mass extinction** is **anthropogenic**: it is caused by human activity, unlike the earlier mass extinctions. Brief case studies show a range of causes.

North Island giant moa: loss of terrestrial megafauna

The North Island giant moa (*Dinornis novaeseelandiae*) was a very large flightless bird of New Zealand, where the only native land mammals were bats. It became extinct within about two centuries of the arrival of Polynesian settlers, the ancestors of Māori, around 1300 CE.

- Hunting for meat, and collection of eggs, by people.
- Burning and clearing of forest habitat.
- Slow breeding: small clutches and many years to reach maturity, so populations could not recover from hunting.
- No evolved fear of land predators, so birds were easy to kill.

Caribbean monk seal: loss of a marine species

The Caribbean monk seal (*Neomonachus tropicalis*) lived on beaches, islands and reefs of the Caribbean Sea and Gulf of Mexico. The last confirmed sighting was in 1952, and it was declared extinct in 2008.

- From the late 15th century, European settlers hunted it for oil from its blubber, and for meat and skins. Seals resting on beaches were easy to kill.
- Overfishing of reef fish and invertebrates reduced its food.
- Disturbance of the beaches where it rested and bred.

A local case study: the Arabian ostrich

The Arabian ostrich (*Struthio camelus syriacus*), a subspecies of ostrich, lived in the deserts of the Arabian Peninsula and nearby regions. It was hunted for meat, feathers and eggs for centuries, and declined rapidly in the 20th century once firearms and motor vehicles made hunting far easier. The last reliable records are from around 1939 to 1941, and a single dying bird reported in Jordan in 1966 is the date usually given for its extinction. Any species lost from an area you know can be used as the third case study.


Ecosystem loss

An ecosystem is lost when its community and abiotic conditions are changed so much that it no longer functions as before. Only causes that are directly or indirectly **anthropogenic** are studied here.

Mixed dipterocarp forest in Southeast Asia. These lowland rainforests of Borneo, Sumatra and the Malay Peninsula are dominated by tall trees of the family Dipterocarpaceae and are home to very many species, including orangutans. Large areas have been lost.

- Logging of valuable dipterocarp hardwood timber.
- Clearance of land for oil palm plantations, and for rubber and pulpwood.
- Fires set to clear land, which spread further in drought years and on drained peat soils.
- Roads and mining that open up remaining forest, leaving small fragments too small for many species.

A local case: coral reefs of the southern Arabian Gulf. Reefs off the UAE already live close to the upper temperature limit for corals. In summer 2017, weak winds failed to cool the shallow sea, and water stayed hot enough for long enough to bleach about 95 % of the corals off Abu Dhabi. By April 2018 about three-quarters (73 %) of them had died, including normally heat-tolerant species. The causes are indirectly anthropogenic, as climate change makes marine heatwaves more frequent, and directly anthropogenic, as dredging, land reclamation for coastal development and sewage discharge add stress. Another well-known example is the Aral Sea in Central Asia, whose lake ecosystem largely collapsed after its feeder rivers were diverted to irrigate cotton from the 1960s.



[\(opens full size in a new tab\)](#)

Anthropogenic causes and effects of the **loss of mixed dipterocarp forest** in Southeast Asia.

Exam tip: Give the specific cause for each case study: overhunting of a slow-breeding bird for the moa, hunting for oil and loss of food for the monk seal, and logging, plantations and fire for dipterocarp forest.

Evidence for and causes of the biodiversity crisis Spec A4.2.5, A4.2.6

Evidence for a crisis

The **Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES)** assesses published evidence on the state of biodiversity. Its 2019 global assessment concluded that around one million animal and plant species are threatened with extinction, and that the current global rate of extinction is already at least tens to hundreds of times higher than the average over the past ten million years.

Such conclusions rest on **reliable surveys** of biodiversity in a wide range of habitats around the world. One survey is only a snapshot: surveys must be **repeated** to show change in two measures:

Species richness	The number of different species present in an area.
Species evenness	How similar the population sizes of the different species are. A community dominated by one species has low evenness.

Practical skill: Repeated surveys are only comparable if the method is standardised: same sites, same season and time of day, same sampling effort and same identification methods.

Percentage change from a repeated survey

A butterfly survey on the same transect recorded 240 individuals in 2010 and 150 in 2020. Calculate the percentage change.

1. Change = $150 - 240 = -90$
2. Percentage change = $\text{change} \div \text{original value} \times 100$
3. = $-90 \div 240 \times 100 = -37.5$

Answer: -37.5 % (a decrease of 37.5 %)

Surveys involve both expert scientists and **citizen scientists**. To be verifiable, evidence usually has to come from a published, peer-reviewed source whose methodology can be checked.

Citizen science	
	Detail
Benefits	Many observers cover far more sites and years than scientists alone could; it builds public interest in conservation
Methodological concerns	Species may be misidentified; effort and methods vary between observers; records cluster where people live

Causes of the current crisis

Human population growth is the overarching cause. More people need more land, food, water, energy and materials. It acts through these specific causes:

Specific causes of biodiversity loss		
Cause	How it reduces biodiversity	Example
Hunting and other over-exploitation	Removes individuals faster than populations can reproduce	Overfishing of cod; hunting of the moa
Urbanization	Replaces natural habitat with buildings and roads, and fragments what is left	Coastal development replacing mangroves
Deforestation and clearance of land for agriculture	Loss of natural habitat	Dipterocarp forest cleared for oil palm
Pollution	Kills organisms or alters habitats	Fertiliser run-off causing eutrophication; plastics in the oceans
Spread of pests, diseases and invasive alien species by global transport	New species outcompete, prey on or infect native species with no defences	Chytrid fungus in amphibians; brown tree snakes eliminating forest birds on Guam

Exam tip: Linking question: how does variation contribute to the stability of ecological communities? Communities with more species and more genetic diversity are more likely to keep functioning when conditions change.

Approaches to conservation and the EDGE of Existence programme Spec A4.2.7, A4.2.8

No single approach to conservation is sufficient by itself, and **different species require different measures**. A species with a large range needs protected habitat; one reduced to a few individuals may need captive breeding.


Approaches to conservation		
Approach	What it involves	Strength or limitation
In situ conservation	Protecting species in their natural habitats	Keeps species interacting with their community and still evolving; fails if the habitat is too degraded
Management of nature reserves	Protecting and actively managing land or sea: controlling invasive species, preventing poaching, managing grazing and fire, linking reserves with corridors	Needs long-term funding and enforcement
Rewilding	Reintroducing lost species, often large herbivores or top predators, and letting natural processes restore the ecosystem	Wolves reintroduced to Yellowstone National Park in 1995 changed grazing by elk
Reclamation of degraded ecosystems	Restoring damaged land or water, for example replanting mangroves or re-flooding drained wetlands	Slow and costly, but recovers whole communities
Ex situ conservation in zoos	Captive breeding of threatened animals, then reintroduction	The Arabian oryx, extinct in the wild by 1972, was bred in captivity and reintroduced from 1982
Botanic gardens	Growing rare plants away from their habitat	Small populations have low genetic diversity
Seed and tissue banks	Storing germ plasm: dried seeds kept cold, or tissue samples frozen	Stores huge genetic diversity in a small space, e.g. the Svalbard Global Seed Vault for crop seeds

The EDGE of Existence programme

Conservation money and effort are limited, so species must be prioritised. The EDGE of Existence programme, run by the Zoological Society of London, focuses on species that are both **Evolutionarily Distinct** and **Globally Endangered**.

- **Evolutionarily distinct (ED):** the species has few or no close living relatives, so it sits at the end of a long branch of the evolutionary tree and represents a lot of unique evolutionary history.
- **Globally endangered (GE):** the species is at high risk of extinction, judged by its IUCN Red List category.
- A species scoring high on both is an EDGE species and a high priority.

The rationale: if a species with no close relatives becomes extinct, a disproportionate amount of unique genes, features and evolutionary history is lost, and no similar species remains. Many EDGE species, such as the aye-aye of Madagascar, the Chinese giant salamander, pangolins and the gharial, receive little attention from other programmes.



[\(opens full size in a new tab\)](#)

An **EDGE** species is both **evolutionarily distinct** (long branch, few relatives) and **globally endangered**.

Exam tip: Nature of science: deciding which species to prioritise has ethical, environmental, political, social, cultural and economic implications. In discuss questions, give arguments for and against the EDGE approach.

Quick check

1. Name the three levels of biodiversity.
2. What does fossil evidence suggest about the number of species alive today?
3. Why must biodiversity surveys be repeated?
4. What is the overarching cause of the current biodiversity crisis?
5. What does EDGE stand for?

Quick check answers

Evolution and speciation

1. Change in the heritable characteristics of a population.
2. They shared a common ancestor more recently.
3. Humerus; radius and ulna; carpals; metacarpals; phalanges.
4. Reproductive isolation and differential selection.

Conservation of biodiversity

1. Ecosystem diversity, species diversity and genetic diversity.
2. There are more species alive today than at any time in the past.
3. To provide evidence of change in species richness and evenness over time.
4. Human population growth.
5. Evolutionarily distinct and globally endangered.