



Proteins: practice questions

4 questions • 11 marks • mark scheme at the back

For IB Diploma Biology SL undefined. **HL only** is additional higher level content, for HL students only.

Name

Total / 11

1 PAPER 1A STYLE [1 mark]

Between which groups does a peptide bond form when two amino acids are joined?

- A. ☐ The R-group of one amino acid and the R-group of the other
- B. ☐ The carboxyl group of one amino acid and the amine group of the other
- C. ☐ The amine groups of both amino acids
- D. ☐ The alpha carbon of one amino acid and the hydrogen of the other

2 PAPER 1A STYLE [1 mark]

How many different tripeptides could be made from the 20 amino acids coded for in the genetic code?

- A. ☐ 60
- B. ☐ 400
- C. ☐ 1140
- D. ☐ 8000

3 PAPER 1B STYLE

[6 marks]

Samples of an enzyme from a mammal were kept at different temperatures for 10 minutes, then cooled to 37 °C. The activity of each sample was then measured at 37 °C and given as a percentage of the activity of an unheated sample. The data are for practice. (a) Describe the effect of pre-incubation temperature on enzyme activity. [2] (b) Calculate the percentage decrease in activity between 50 °C and 60 °C. [1] (c) Explain why the activity of the samples kept at 70 °C and 80 °C did not recover when cooled. [2] (d) Suggest why activity was measured at 37 °C for every sample. [1]

Activity after 10 minutes at each temperature (practice data)

Pre-incubation temperature / °C	Activity / % of unheated sample
20	100
40	98
50	80
60	35
70	6
80	0

.....

.....

.....

.....

.....

.....

.....

.....

4 PAPER 2A STYLE

[3 marks]

Distinguish between essential and non-essential amino acids and outline why people following a vegan diet need to pay attention to protein sources.

.....

.....

.....

.....

.....

Mark scheme

One mark for each point unless a maximum is given. Words in brackets are not needed for the mark.

1. [1]

B B [1]

2. [1]

D D [1]

3. [6]

- (a) little / no loss of activity up to 40 °C [1]
- (a) above 40 °C activity falls, steeply between 50 °C and 70 °C / no activity at 80 °C [1]
- (b) $(80 - 35) \div 80 \times 100 = 56\%$ [1] Accept 56.25 %
- (c) the enzyme was denatured / its three-dimensional shape / active site changed [1]
- (c) hydrogen bonds / weak bonds broken by increased vibration; denaturation is (usually) permanent so cooling does not restore the shape [1]
- (d) so that any difference in activity is due only to the effect of the pre-incubation / irreversible change / to control temperature during the assay [1]

4. [3]

- essential amino acids cannot be synthesised by the body / must be obtained from food [1]
- non-essential amino acids can be made from other amino acids [1]
- some plant proteins contain little of certain essential amino acids [1]
- so vegans need a variety of plant protein sources / combine e.g. legumes and cereals [1]
- max 3